


Cultural eutrophication in the Greater North Sea

Cause, symptoms, mitigation



How can science guide ecologically-relevant and economically sustainable decisions ?

Christiane Lancelot
Ecologie des Systèmes Aquatiques

CONTENT

I-The Global Eutrophication Context

II-Eutrophication in the Greater North Sea


III-Science Support to Combat Eutrophication

I-The Global Eutrophication Context

- ✓ Global trends in nutrient loads
- ✓ Mechanisms behind eutrophication
- ✓ Undesirable effects

Global Nutrient Loads

Since 1950: the Great Acceleration: a connected global system
population * socio-economy * green revolution



As a consequence:

Global increase of N (18%), P (13%) river inputs

Seitzinger et al. 2010

Global retention (18%) of Si

Beusen et al., 2009

Exacerbated by release from growing aquaculture, atmospheric and groundwater inputs

Cultural Eutrophication

Modification of the natural N:P:Si balance of coastal waters towards N and P excess → primary production ↗

Undesirable if this nutrient excess appreciably degrades ecosystem health and/or the provision of goods and services

MSD-TG5 report 2010; Ferreira et al. 2011

DESIRABLE

Interannual variations



Nutrient loads
Total N, P: ↗
Si: ≈ or ↘

Change in coastal
N:P:Si nutrient status

UNDESIRABLE




HABs (high-biomass, toxic...)
Hypoxia/anoxia
Benthic/pelagic species shifts
...

Different symptoms !


The functioning of Marine Ecosystems




Marine Ecosystem: Generic Process





Pristine
N:Si:P=16:16:1
phytoplankton pattern
Dominated by DA



Cultural eutrophication: Generic Process



Eutrophied phytoplankton pattern



Undesirable symptoms of eutrophication: HABs



Foam deposits




Algal deposit & H₂S release



Fish mortality

Shellfish poisoning

High-biomass HABs




Phaeocystis globosa



Ulva

Toxic HABs



Karenia mikimotoi




Pseudo-Nitzschia



Dinophysis

II-Eutrophication in the Greater North Sea



- ✓ Historical key dates
- ✓ Nutrient sources today
- ✓ Eutrophication symptoms

Eutrophication in the greater North Sea: key dates

✓ 1950-1970

Great Acceleration

✓ After 1970

Eutrophication symptoms

Environmental awareness

✓ 1975-1991

Governance (OSPAR, HELCOM)

Combination of professional judgment and political art :e.g. 50% ↘ 1985 N and P loads




First EU Directives (WWT, Nitrate)

✓ Since 2000 (WFD and MSD)

Awareness of the 'land-sea' connectivity, regime shifts, thresholds, points of non-return → Ecological Quality Objectives.

Sustainable coastal sea.


Nutrient river load
e.g. River Scheldt



Model reconstruction after Lancelot et al., 2007;
Passy et al., 2013


Eutrophication in the greater North Sea: current status

Ocean color 2011




Nutrient enrichment

Winter nitrate concentration



Atmospheric deposition
fraction total N






Same causes but distinct blooms

III-Science Support to Combat Eutrophication

What can be done to improve the situation and how can we appraise mitigation actions ?

- ✓ Integrated Impact Assessment Pathway/toolkit
- ✓ Case study: *Phaeocystis* blooms

Impact Assessment Pathway: integration of coupled biogeochemical models in a DPSIR loop








Lancelot et al., 2011

(1) Ruelland et al. 2007; (2) Billen and Garnier 1999; (3) Lancelot et al., 2005

Some possible scenarios

Garnier et al., 2012



Overall conclusion and futuring

The fact


Human activity is modifying the quantity and quality of nutrients (N, P, Si) delivered to the coastal sea;

This N & P enrichment has boosted non-diatom species often undesirable/harmful;

Most attempts to reducing N & P loads have not decrease coastal problems because of end-of-pipe solution choice rather than acting on the NUE (nutrient use efficiency) process.

Towards solutions: de-eutrophication toolkit

- ✓ Interlinked suite of ecological and socio-economic models
- ✓ Scenario-driven approach: dialogue between science and society



Acknowledgements



EU FP6

Thresholds of Environmental sustainability

Belgian Science Policy Office



belspo

IAP TIMOTHY (**T**racing and **I**ntegrated
Modeling of Natural and Anthropogenic
Effects on **H**ydrosystems (TIMOTHY)
Case study: The Scheldt River Basin and
Adjacent Coastal North Sea

AMORE (**A**dvanced **M**odelling and **R**esearch
on **E**utrophication)



EU FP7

AWARE

Adaptative management: Increased
connectivity between politic, science, public



Ecosystem **M**odels as **S**upport to **E**utrophication
Management In the North Atlantic Ocean

Cited references

- Beusen, A.H.W., Bouwmann, A.F., Durr, H.H., Dekkers, A.L.M., Hartmann, J. Global patterns of dissolved silica export to the coastal zone: results from a spatially explicit global model. *Global Biogeochem. Cycles*, **23**:GB0A02, doi: 10.1029/2008GB003281 (2009)
- Billen, G., Garnier, J., 1999. Nitrogen transfers through the Seine drainage network: a budget based on the application of the Riverstrahler model. *Hydrobiol.* 410, 139-150
- Ferreira JG, Andersen JH, Borja A, et al. (2011) Overview of eutrophication indicators to assess environmental status within the European Marine Strategy Framework Directive RID C-5701-2011. *Estuar Coast Shelf Sci* 93:117–131. doi: 10.1016/j.ecss.2011.03.014.
- Garnier J, Passy P, Rousseau V, Gypens N, Callens J, Silvestre M, Billen J, Lancelot C. 2012. The diseased southern North Sea: current status and possible solutions. In: Sustainable Water Ecosystems Management in Europe. Bridging the knowledge of citizens, scientists and policy-makers. Ed C. Sessa. EU report series, IWA Publishing, London, NY.
- Lancelot C., Spitz, Y., Gypens, N., Ruddick, K., Becquevort, S., Rousseau, V., Lacroix, G., Billen, G., 2005. Modelling diatom and *Phaeocystis* blooms and nutrient cycles in the Southern Bight of the North Sea: the MIRO model. *Mar. Ecol. Progr. Ser.* 289, 63-78.
- Lancelot, C., Gypens, N., Billen, G., Garnier, J., Roubex, V., 2007. Testing an integrated river-ocean mathematical tool for linking marine eutrophication to land use: the *Phaeocystis*-dominated Belgian coastal zone (Southern North Sea) over the past 50 years. *J. Mar. Sys.* 64, 216–228.
- Lancelot, C., Rousseau, V., Gypens, N., 2009. Ecologically based indicators for *Phaeocystis* disturbance in eutrophied Belgian coastal waters (Southern North Sea) based on field observations and ecological modelling. *J. Sea Res.* 61, 44-49.
- Lancelot, C., Thieu, V., Polard, A., Garnier, J., Billen, G., Hecq, W., Gypens, N., 2011. Cost assessment and ecological effectiveness of nutrient reduction options for mitigating *Phaeocystis* colony blooms in the Southern North Sea: An integrated modeling approach. *Sci. Total Environ.* 409, 2179–2191.
- Lancelo C, Passy P, Gypens N. 2014. Model assessment of present-day *Phaeocystis* colony blooms in the Southern Bight of the North Sea (SBNS) by comparison with a reconstructed pristine situation. *Harmful Algae* 37:172-182.
- Passy, P., Gypens, N., Billen, G., Garnier, J., Lancelot, C., Thieu, V., Rousseau, V., Callens, J., 2013. A Model reconstruction of riverine nutrient fluxes and eutrophication in the Belgian Coastal Zone since 1984, 2013. *J. Mar. Syst.* 128, 106-122.
- Ruelland, D., Billen, G., Brunstein, D., Garnier, J., 2007. SENEQUE 3: a GIS interface to the RIVERSTRAHLER model of the biogeochemical functioning of river systems. *Sci. Tot. Environ.* 375, 257-273.
- Seitzinger, S.P. *et al.* Global river nutrient export: a scenario analysis of past and future trends. *Global Biogeochem. Cycles*, **24**: GB0A08, doi:10.1029/2009GB003587 (2010).
- Sivystski, J. 2012. Anthropocene: an epoch of our making. *Global Change* issue 78.
- Troost, TA, Blaas M, LOS FJ. 2013. The role of atmospheric deposition in the eutrophication of the North Sea: a model analysis. *J. Mar. Sys.* 125: 101-112